

## SCIENTIFIC ARTICLES

© A. Ponosov, 2023

DOI 10.20310/2686-9667-2023-28-141-51-59



## Existence and uniqueness of solutions to stochastic fractional differential equations in multiple time scales

Arcady PONOSOV

Norwegian University of Life Sciences  
№-1432 Ås 5003, Drøbakveien 31, Norway

**Abstract.** A novel class of nonlinear stochastic fractional differential equations with delay and the Jumarie and Itô differentials is introduced in the paper. The aim of the study is to prove existence and uniqueness of solutions to these equations. The main results of the paper generalise some previous findings made for the non-delay and three-scale equations under additional restrictions on the fractional order of the Jumarie differentials, which are removed in our analysis. The techniques used in the paper are based on the properties of the singular integral operators in specially designed spaces of stochastic processes, the representation of delay equations as functional differential equations as well as Picard's iterative method.

**Keywords:** Jumarie derivative, Brownian motion, multi-time scales

**Mathematics Subject Classification:** 60H20, 34A8, 34K50.

**Acknowledgements:** The author gratefully acknowledges the financial support from internal funding scheme at Norwegian University of Life Sciences (project no. 1211130114), which financed the international stay at Scuola Normale Superiore in Italy.

**For citation:** Ponosov A. Existence and uniqueness of solutions to stochastic fractional differential equations in multiple time scales. *Vestnik rossiyskikh universitetov. Matematika = Russian Universities Reports. Mathematics*, 2023, vol. 28, no. 141, pp. 51–59. DOI 10.20310/2686-9667-2023-28-141-51-59.

НАУЧНАЯ СТАТЬЯ

© Поносков А.В., 2023

DOI 10.20310/2686-9667-2023-28-141-51-59

УДК 519.216.2



## Существование и единственность решений стохастических дробных дифференциальных уравнений в нескольких временных шкалах

Аркадий Владимирович ПОНОСОВ

Норвежский университет естественных наук

1432, Норвегия, г. Ос, Дрёбаквайен, 31

**Аннотация.** В статье вводится новый класс нелинейных стохастических дифференциальных уравнений дробного порядка с запаздыванием и дифференциалами Жюмари и Итô. Цель исследования — доказать существование и единственность решений этих уравнений. Основные результаты статьи обобщают некоторые предыдущие выводы, сделанные для уравнений без запаздывания с тремя временными шкалами и при дополнительных ограничениях на дробный порядок дифференциалов Жюмари, которые снимаются в нашем анализе. Методы, использованные в статье, основаны на свойствах сингулярных интегральных операторов в специально сконструированных пространствах случайных процессов, представлении уравнений с запаздыванием в виде функционально-дифференциальных уравнений, а также на итерационном методе Пикара.

**Ключевые слова:** производная Жюмари, броуновское движение, мультивременные шкалы

**Благодарности:** Автор выражает благодарность за финансовую поддержку работы в рамках внутренней программы Норвежского университета естественных наук (проект № 1211130114), которая финансировала международный визит в Высшую нормальную школу в Италии.

**Для цитирования:** Поносков А.В. Существование и единственность решений стохастических дробных дифференциальных уравнений в нескольких временных шкалах // Вестник российских университетов. Математика. 2023. Т. 28. № 141. С. 51–59. DOI 10.20310/2686-9667-2023-28-141-51-59. (In Engl., Abstr. in Russian)

## 1. Introduction.

Processes operating in a multi-time scale modus arise in a number of fields including finance, science and engineering. In [1] it was suggested to use the fractional Jumarie derivative introduced in [2]  $v = \frac{df}{(dt)^\alpha}$  ( $0 < \alpha \leq 1$ ) and the classical white noise  $g = w \frac{dB}{dt}$  to model the deterministic and the stochastic parts of the multi-time scale processes, respectively. In the integral form this reads as

$$f(t) - f(0) = \alpha \int_0^t (t-s)^{\alpha-1} v(s) ds \quad \text{and} \quad g(t) - g(0) = \int_0^t w(s) dB(s).$$

Adopting this approach we study the following fractional stochastic delay differential equation in multiple time scales:

$$dx(t) = \sum_{j=1}^m (f_j(t, (H_{1j}x)(t))(dt)^{\alpha_j} + g_j(t, (H_{2j}x)(t))dB_j(t)). \quad (1.1)$$

Here  $f_j(t, v)$  and  $g_j(t, v)$  are random functions and  $H_{1j}$ ,  $H_{2j}$  are linear delay operators,  $(dt)^{\alpha_j}$  are the fractional Jumarie differentials and  $dB_j(t)$  are the Itô differentials generated by the standard scalar Wiener processes (Brownian motions)  $B_j$ . The initial condition for (1.1) is

$$x(s) = \varphi(s) \quad (s \leq 0), \quad (1.2)$$

where  $\varphi(\omega, s)$  is some random function (not necessarily continuous).

A solution of the initial value problem (1.1), (1.2) is a stochastic process  $x$  satisfying (1.2) for  $s \leq 0$  and the integral equation

$$x(t) - \varphi(0) = \sum_{j=1}^m \left( \int_0^t \alpha_j (t-s)^{\alpha_j-1} f_j(s, (H_{1j}x)(s)) ds + \int_0^t g_j(s, (H_{2j}x)(s)) dB_j(s) \right).$$

The main result of the paper is a generalization of the existence and uniqueness theorem from [1] to the case of Eq. (1.1) and its operator counterpart.

## 2. Preliminaries

We keep fixed a stochastic basis  $(\Omega, \mathcal{F}, (\mathcal{F})_{t \in R}, P)$  satisfying the standard conditions [3] assuming, in addition, that  $\mathcal{F}_t = \mathcal{F}_0$  for all  $t \leq 0$ . All stochastic processes in this paper are supposed to be progressively measurable w.r.t. this stochastic basis or parts of it [3].

The following notation is used throughout the paper:

- $R = (-\infty, \infty)$ ,  $R_+ = [0, \infty)$ ,  $R_- = (-\infty, 0)$ .
- $\mu$  is the Lebesgue measure defined on  $R$  or its subintervals.
- $E$  is the expectation corresponding to the probability measure  $P$ .
- $B_j(t)$  ( $t \in R_+$ ,  $j = 1, \dots, m$ ) are the standard scalar Wiener processes.
- The space  $\mathcal{L}_q(J, R^l)$  ( $1 \leq q < \infty$ ,  $J \subset R$  is a subinterval), contains all progressively measurable  $l$ -dimensional stochastic processes  $x(t)$  ( $t \in J$ ) such that  $\int_J E|x(t)|^q dt < \infty$ .
- The space  $\mathcal{M}_p(J, R^l)$  ( $1 \leq p < \infty$ ) consists of all progressively measurable,  $l$ -dimensional stochastic processes  $x(t)$  ( $t \in J$ ) such that

$$\sup_{t \in J} E|x(t)|^p < \infty.$$

- The space  $k^n$  consists of all  $n$ -dimensional,  $\mathcal{F}_0$ -measurable random variables, and  $k = k^1$  is a commutative ring of all scalar  $\mathcal{F}_0$ -measurable random variables.

- The space  $k_p^n = \{\xi : \xi \in k^n, E|\xi|^p < \infty\}$  ( $1 \leq p < \infty$ ) is a linear subspace of  $k^n$ .

The spaces  $\mathcal{L}_q(J, R^l)$ ,  $\mathcal{M}_p(J, R^l)$  and  $k_p^n$  are supposed to be equipped with the natural norms. Clearly also that for  $q \leq p$  and finite intervals  $J$  we have  $\mathcal{M}_p(J, R^l) \subset \mathcal{L}_q(J, R^l)$ , also in the topological sense (the topology of the larger space is weaker). We will also assume that  $\mathcal{M}_p(J, R^l) \subset \mathcal{M}_p(J', R^l)$  if  $J \subset J'$  by letting the processes on  $J$  to be 0 outside  $J$ .

### 3. Properties of some delay operators

Consider the delay operator

$$(Hx)(t) = x(h(t)), \quad (3.1)$$

**Theorem 3.1.** *Let  $J = [0, T]$  and  $1 \leq q < \infty$ . Assume that  $h(t)$  ( $t \in J$ ) is a Borel function such that  $h(t) \leq t$   $\mu$ -almost everywhere on  $J$ . Then the operator (3.1) is a linear bounded operator from  $\mathcal{M}_q(R_- \cup J, R^n)$  to  $\mathcal{M}_q(J, R^n)$ .*

*P r o o f.* Evidently,  $H$  is linear and maps progressively measurable processes defined on  $R_- \cup J$  to the ones defined on  $J$ . In addition,  $\sup_{t \in J} E|x(h(t))|^q \leq \sup_{t \leq T} E|x(t)|^q$ , which proves boundedness of  $H$  from  $\mathcal{M}_q(R_- \cup J, R^n)$  to  $\mathcal{M}_q(J, R^n)$ .  $\square$

Next, consider the distributed delay operator

$$(Hx)(t) = \int_{-\infty}^t d_s \mathcal{R}(t, s)x(s). \quad (3.2)$$

**Theorem 3.2.** *Let  $J = [0, T]$  and  $1 < q < \infty$ . Assume that the values of  $\mathcal{R}(t, s)$  ( $t \in J, -\infty < s \leq t$ ) are  $l \times n$ -matrices and  $\mathcal{R}$  satisfies the following conditions:*

1.  $\mathcal{R}$  is Borel measurable on its domain;
2.  $\sup_{t \in J} \text{Var}_{-\infty}^t \mathcal{R}(t, \cdot) < \infty$ .

*Then the operator (3.2) is a linear bounded operator from  $\mathcal{M}_r(R_- \cup J)$  to  $\mathcal{M}_q(J, R^l)$ .*

*P r o o f.* Using the componentwise description of the operator (3.2) we may assume, without loss of generality, that  $l = n = 1$ , so that  $(Hx)(t) = \int_{-\infty}^t x(s)d_s \mathcal{R}(t, s)$ . Evidently, the operator  $H$  maps progressively measurable processes defined on  $R_- \cup J$  to the ones defined on  $J$ . Putting  $\text{Var}_{-\infty}^t[\mathcal{R}(t, \cdot)](s) \equiv \bar{\mathcal{R}}(t, s)$  we get

$$\begin{aligned} & \sup_{t \in J} E \left| \int_{-\infty}^t x(s)d_s \mathcal{R}(t, s) \right|^q \\ & \leq \sup_{t \in J} \left( E \int_{-\infty}^t |x(s)|^q d_s \bar{\mathcal{R}}(t, s) \times \left( \int_{-\infty}^t d_s \bar{\mathcal{R}}(t, s) \right)^{q-1} \right) \\ & \leq \sup_{t \in J} \left( \int_{-\infty}^t E|x(s)|^q d_s \bar{\mathcal{R}}(t, s) \right) \times \sup_{t \in J} \left( \int_{-\infty}^t d_s \bar{\mathcal{R}}(t, s) \right)^{q-1} \\ & \leq \sup_{s \in R_- \cup J} E|x(s)|^q \sup_{t \in J} \left( \int_{-\infty}^t d_s \bar{\mathcal{R}}(t, s) \right) \times \sup_{t \in J} \left( \int_{-\infty}^t d_s \bar{\mathcal{R}}(t, s) \right)^{q-1} \\ & \leq \sup_{t \in R_- \cup J} E|x(t)|^q \sup_{t \in J} \left( \int_{-\infty}^t d_s \bar{\mathcal{R}}(t, s) \right)^q \leq \left( \sup_{t \in J} \text{Var}_{-\infty}^t \mathcal{R}(t, \cdot) \right) \sup_{t \in R_- \cup J} E|x(t)|^q \end{aligned}$$

which proves boundedness of  $H$  from  $\mathcal{M}_q(R_- \cup J)$  to  $\mathcal{M}_q(J, R^l)$ .  $\square$

**R e m a r k 3.1.** The delay operator (3.1) can be regarded as a particular case of the delay operator (3.2) if one puts  $\mathcal{R}(t, s) = \text{diag}[\chi_h, \dots, \chi_h]$  to be the  $n \times n$  diagonal matrix containing the indicator  $\chi_h$  of the set  $\{(t, s) : s \leq h(t)\}$ . Moreover, if we define  $\mathcal{R}(t, s)$  to be the  $(rn) \times n$ -matrix of the form

$$\mathcal{R}(t, s) = (\text{diag}[\chi_{h_1}, \dots, \chi_{h_1}], \dots, \text{diag}[\chi_{h_r}, \dots, \chi_{h_r}]),$$

then we get the multiple delay operator  $x(t) \mapsto (x(h_1(t)), \dots, x(h_r(t)))$ .

#### 4. Main results

Let us first consider the following fractional functional differential equation:

$$dy(t) = \sum_{j=1}^m ((F_j(y))(t)(dt)^{\alpha_j} + (G_j(y)x)(t)dB_j(t)), \quad t \in [0, T] \quad (4.1)$$

equipped with the initial condition

$$y(0) = y_0 \in k_p^n. \quad (4.2)$$

The solution of Eq. (4.1) is understood in the following sense:

$$y(t) - y(0) = \sum_{j=1}^m \left( \alpha_j \int_0^t (t-s)^{\alpha_j-1} F_j(y)(s) ds + \int_0^t G_j(y)(s) dB_j(s) \right), \quad t \in [0, T]. \quad (4.3)$$

**D e f i n i t i o n 4.1.** Let  $X, Y$  be two separable metric spaces and  $\sigma : \Omega \times X \rightarrow Y$  be a random map. The operator  $x(\cdot) \rightarrow \sigma(\cdot, x(\cdot))$  is called the superposition operator generated by  $\sigma$ .

**D e f i n i t i o n 4.2.**

- A continuous map  $V : X \rightarrow Y$ , where  $X, Y$  are two separable metric spaces of functions defined on an interval  $J \subset R$  is called Volterra if

$$x_1(s) = x_2(s) \quad \Rightarrow \quad (Vx)(s) = (Vy)(s)$$

for all  $x_1, x_2 \in \mathcal{X}$ , any  $t \in J$  and almost all  $s \leq t, s \in J$ .

- A map  $V : \Omega \times X \rightarrow Y$  is called a random Volterra map if  $V(\omega, \cdot)$  is Volterra for almost all  $\omega \in \Omega$  and  $V(\cdot, x)$  is  $\mathcal{F}$ -measurable for all  $x \in X$ .
- The superposition operator generated by a random Volterra map is defined by  $x(\cdot) \mapsto V(\cdot, x(\cdot))$ .
- A random Volterra map  $V : \Omega \times X \rightarrow Y$ , such that  $V^t(\cdot, x)$  is  $\mathcal{F}_t$ -measurable for all  $t \in J$  will be called non-anticipating.

Evidently, any Volterra map  $V$  gives rise to a family of Volterra maps  $V^t : X_t \rightarrow Y_t$  ( $t \in J$ ), where  $X_t$  and  $Y_t$  consist of the restrictions of the functions from  $X$  and  $Y$ , respectively, to  $(-\infty, t] \cap J$ . It is also easy to check that the superposition operators generated by random Volterra maps are continuous in probability and if  $V$  is non-anticipating, then the superposition operator generated by  $V$  preserves progressive measurability of stochastic processes.

In the proofs below we use the following inequalities:

$$E \left| \int_0^t f(s) dB(s) \right|^q \leq c_q^q E \left( \int_0^t |f(s)|^2 ds \right)^{q/2} \quad (t \in R_+, q \geq 2), \quad (4.4)$$

where  $f(t)$  is an arbitrary scalar, progressive measurable stochastic process on  $R_+$ ,  $B(t)$  is the standard scalar Brownian motion and  $c_q$  is a certain constant, which is independent of  $f$ ;

$$\left| \int_0^t (t-s)^{\alpha-1} g(s) ds \right|^q \leq d_q^q t^{q\alpha-1} \int_0^t |g(s)|^q ds \quad (t \in R_+, q > \alpha^{-1}), \quad (4.5)$$

where  $g : R_+ \rightarrow R$  is a Lebesgue measurable function and  $d_q = \left( \frac{q-1}{q\alpha-1} \right)^{1-1/q}$ .

Inequality (4.4) follows from the estimates proved in e.g. [4], while (4.5) is a direct consequence of Hölder's inequality.

**Theorem 4.1.** *Let  $J = [0, T]$  and assume that*

1.  $0 < \alpha_j \leq 1$ ,  $p_j \geq 2$ ,  $\alpha_j^{-1} < p_j \leq p$  ( $1 \leq j \leq m$ ).
2. *The superposition operators generated by the non-anticipating operators  $F_j, G_j$  ( $1 \leq j \leq m$ ) map the space  $\mathcal{M}_p(J, R^n)$  into the spaces  $\mathcal{L}_{p_j}(J, R^n)$  and  $\mathcal{L}_2(J, R^n)$ , respectively, and satisfy the Lipschitz condition*

$$\begin{aligned} \|F_j y_1 - F_j y_2\|_{\mathcal{L}_{p_j}(J, R^n)} &\leq \ell \|y_1 - y_2\|_{\mathcal{M}_p(J, R^n)}, \\ \|G_j y_1 - G_j y_2\|_{\mathcal{L}_2(J, R^n)} &\leq \ell \|y_1 - y_2\|_{\mathcal{M}_p(J, R^n)} \end{aligned} \quad (4.6)$$

for some constant  $\ell$  and the sub-linear growth condition

$$\|F_j \xi\|_{\mathcal{L}_{p_j}(J, R^n)} \leq b \|\xi\|_{k_p^n}, \quad \|G_j \xi\|_{\mathcal{L}_2(J, R^n)} \leq b \|\xi\|_{k_p^n} \quad (4.7)$$

for some constant  $b$  and any  $\xi \in k_p^n$ .

Then the initial value problem (4.1), (4.2) has a unique (up to the natural equivalence of indistinguishable processes) solution  $y(\cdot, y_0) \in \mathcal{M}_p(J, R^n)$ .

If the constants  $\ell$  and  $b$  are independent of  $J$ , then the solution  $y(t, y_0)$  is defined for all  $t \in R_+$ .

**P r o o f.** We prove this theorem for the equivalent integral equation (4.3). Notice that due to the Volterra property of the operators  $F_j$  and  $G_j$  we have

$$\begin{aligned} \|F_j y_1 - F_j y_2\|_{\mathcal{L}_{p_j}([0, t], R^n)} &\leq \ell \|y_1 - y_2\|_{\mathcal{M}_p([0, t], R^n)}, \\ \|G_j y_1 - G_j y_2\|_{\mathcal{L}_2([0, t], R^n)} &\leq \ell \|y_1 - y_2\|_{\mathcal{M}_p([0, t], R^n)} \end{aligned} \quad (4.8)$$

for any  $t \in J$ . Now, the proof becomes a standard application of Picard's iterations. Put

$$y^{(\nu)}(t) = y_0 + \sum_{j=1}^m \left( \alpha_j \int_0^t (t-s)^{\alpha_j-1} F_j(y^{(\nu-1)})(s) ds + \int_0^t G_j(y^{(\nu-1)})(s) dB_j(s) \right) \quad (t \in J, \nu \in N)$$

and  $y^{(0)} = y_0$ . Using (4.6)–(4.8) and inequalities (4.4), (4.5) with  $q = p_j$  we obtain

$$E |y^{(\nu+1)}(t) - y^{(\nu)}(t)|^p \leq K \int_0^t E |y^{(\nu)}(s) - y^{(\nu-1)}(s)|^p ds \quad (t \in J, \nu \in N) \quad (4.9)$$

and

$$E |y^{(1)}(t) - y^{(0)}(t)|^p \leq K_0 t \|y_0\|_{k_p^p} \quad (t \in J). \quad (4.10)$$

Iterating (4.9) and using (4.10) yield

$$E |y^{(\nu+1)}(t) - y^{(\nu)}(t)|^p \leq K_0 \frac{K^\nu t^\nu}{\nu!} \quad (t \in J, \nu \in N),$$

which ensures convergence of the sequence  $\{y^{(\nu)}\}$  to some  $y$  in the space  $\mathcal{M}_p(J, R^n)$ . The stochastic process  $y(t)$  satisfies then Eq. (4.3) due to continuity of the operators

$$F_j : \mathcal{M}_p(J, R^n) \rightarrow \mathcal{L}_{p_j}(J, R^n) \quad \text{and} \quad G_j : \mathcal{M}_p(J, R^n) \rightarrow \mathcal{L}_2(J, R^n)$$

and boundedness of the linear operators

$$(\mathcal{I}_{1j}y)(t) = \int_0^t (t-s)^{\alpha_j-1} y(s) ds \quad \text{and} \quad (\mathcal{I}_{2j}y)(t) = \int_0^t y(s) dB(s)$$

acting from  $\mathcal{L}_{p_j}(J, R^n)$  to  $\mathcal{M}_p(J, R^n)$  and from  $\mathcal{L}_2(J, R^n)$  to  $\mathcal{M}_p(J, R^n)$ , respectively (see estimates (4.4), (4.5)).

Assume  $y_1(t)$  and  $y_2(t)$  to be two solutions of Eq. (4.3). Then we have, exactly as in (4.9), that

$$E |y_1(t) - y_2(t)|^p \leq K \int_0^t E |y_1(s) - y_2(s)|^p ds \quad (t \in J),$$

and the property of uniqueness follows from Grönwall's lemma.  $\square$

To prove the existence and uniqueness theorem for (1.1) we represent it as Eq. 4.1. This is a standard procedure in the deterministic theory of functional differential equations [5]. To this end, we assume given two stochastic processes  $y \in \mathcal{M}_p(J, R^n)$  and  $\varphi \in \mathcal{M}_p(R_- \cup \{0\})$ , put

$$y_+(t) = \begin{cases} y(t) & (t \in J) \\ 0 & (t \in R_-) \end{cases} \quad \text{and} \quad \varphi_-(t) = \begin{cases} 0 & (t \in J) \\ \varphi(t) & (t \in R_-) \end{cases}$$

and define

$$F_j(y) = f_j(\cdot, H_{1j}y_+ + H_{1j}\varphi_-), \quad G_j(y) = g_j(\cdot, H_{2j}y_+ + H_{2j}\varphi_-), \quad (4.11)$$

which yields Eq. (4.1).

The result below connects Eq. (1.1) and (4.1).

**Proposition 4.1.** *Let  $J = [0, T]$  and assume that the  $k$ -linear operators  $H_{ij} : \mathcal{M}_p(R_- \cup J, R^n) \rightarrow \mathcal{L}_p(J, R^l)$  are bounded for all  $i = 1, 2, j = 1, \dots, m$ . Then the stochastic process*

$$x(t) = \begin{cases} y(t, \varphi(0)) & (t \in J) \\ \varphi(t) & (t \in R_-) \end{cases} \quad (4.12)$$

*is the solution of the initial value problem (1.1), (1.2) on the interval  $J$  if and only if  $y$  is the solution of the initial value problem (4.1), (4.2) on the same interval.*

*P r o o f.* Let  $y$  be a solution of the problem (4.1), (4.2). Then (4.12) can be rewritten as  $x(t) = y_+(t, \varphi(0)) + \varphi_-(t)$  ( $t \in R_- \cup J$ ), and for all  $t \in J$  we obtain  $x(t) = y(t)$  and  $H_{ij}y_+ + H_{ij}\varphi_- = H_{ij}x$  due to linearity of  $H_{ij}$ . Hence  $x(t)$  satisfies Eq. (1.1). In addition,  $x(t) = \varphi(t)$  for  $t \leq 0$ .

Assume now that  $x$  is a solution of the problem (1.1), (1.2) and put  $y = x|_J$ . Then  $x(t) = y_+(t) + \varphi_-(t)$  ( $t \in R_- \cup J$ ), so that  $H_{ij}x = H_{ij}y_+ + H_{ij}\varphi_-$ , which means that  $y(t)$  satisfies Eq. (4.1) if  $F_j$  and  $G_j$  are defined as in (4.11). By construction,  $y(0) = \varphi(0)$ , and the result follows.  $\square$

**E x a m p l e 4.1.** The representation (4.1) of Eq. (1.1) with the distributed delay operators  $H_{ij}$  given by

$$(H_{ij}x)(t) = \int_{-\infty}^t d_s \mathcal{R}_{ij}(t, s)x(s),$$

where  $\mathcal{R}_{ij}(t, s)$  are  $n \times l$ -matrix valued, Borel measurable functions defined on  $\{(t, s) : t \in J, -\infty < s \leq t\}$ , reads as

$$dy(t) = \sum_{j=1}^m \left( f_j(t, \int_0^t d_s \mathcal{R}_{1j}(t, s)y(s) + u_{1j}(t))(dt)^{\alpha_j} + g_j(t, \int_0^t d_s \mathcal{R}_{2j}(t, s)y(s) + u_{2j}(t))dB_j(t) \right),$$

where  $u_{ij}(t) \equiv (H_{ij}\varphi_-)(t) = \int_{(-\infty, 0)} d_s \mathcal{R}_{ij}(t, s)\varphi(s)$ . In particular, Eq. (1.1) with time-dependent delays given by

$$(H_{ij}x)(t) = x(h_{ij}(t)),$$

where  $h_{ij}(t) \leq t$  are Borel measurable functions ( $i = 1, 2, j = 1, \dots, m$ ), has the following representation:

$$dy(t) = \sum_{j=1}^m (f_j(t, (S_{1j}y)(s) + u_{1j}(t))(dt)^{\alpha_j} + g_j(t, (S_{2j}y)(s) + u_{2j}(t))dB_j(t)),$$

where  $S_{ij}$ , known as inner superposition operators (see e.g. [5]), are defined as

$$(S_{ij}y)(t) = \begin{cases} y(h_{ij}(t)) & (t \in J) \\ 0 & (t \in R_-), \end{cases}$$

and

$$u_{ij}(t) \equiv (H_{ij}\varphi_-)(t) = \begin{cases} 0 & (t \in J) \\ \varphi(h_{ij}(t)) & (t \in R_-). \end{cases}$$

Now we are ready to prove the existence and uniqueness result for Eq. (1.1).

**Theorem 4.2.** *Let  $J = [0, T]$  and assume that*

1.  $0 < \alpha_j \leq 1, p_j \geq 2, \alpha_j^{-1} < p_j \leq p$  ( $1 \leq j \leq m$ ).
2. For all  $j = 1, \dots, m$  the random functions  $f_j, g_j : \Omega \times R_+ \times R^l \rightarrow R^n$  are such that  $f_j(\cdot, \cdot, v)$  and  $g_j(\cdot, \cdot, v)$  are progressively measurable for any  $v \in R^l$  and  $f_j(\omega, t, \cdot)$  and  $g_j(\omega, t, \cdot)$  are continuous for  $P \otimes \mu$ -almost all  $(\omega, t)$ , satisfy the Lipschitz condition

$$|f_j(\omega, t, x_1) - f_j(\omega, t, x_2)| \leq \ell|x_1 - x_2|, \quad |g_j(\omega, t, x_1) - g_j(\omega, t, x_2)| \leq \ell|x_1 - x_2| \quad a. s.$$

for some constant  $\ell$  and all  $x_1, x_2 \in R^l, t \in J$  and the sub-linear growth condition

$$|f_j(\omega, t, x)| \leq b|x|, \quad |g_j(\omega, t, x)| \leq b|x|.$$



3. The  $k$ -linear operators  $H_{ij} : \mathcal{M}_p(R_- \cup J, R^n) \rightarrow \mathcal{L}_p(J, R^l)$  are bounded for all  $i = 1, 2, j = 1, \dots, m$ .

Then for any  $\varphi \in \mathcal{M}_p(R_- \cup \{0\}, R^n)$  the initial value problem (1.1), (1.2) has a unique (up to the natural equivalence of indistinguishable processes) solution  $x(\cdot, \varphi) \in \mathcal{M}_p(J, R^n)$ .

If the constant  $\ell$  is independent of  $J$ , then the solution  $x(t, \varphi)$  is defined for all  $t \in R$ .

**P r o o f.** The proof is based on Theorem 4.1. Define  $F_j$  and  $G_j$  using the formulas (4.11). It is easy to see that the superposition operators generated by the non-anticipating operators  $F_j, G_j$  ( $1 \leq j \leq m$ ) map the space  $\mathcal{M}_p(J, R^n)$  into the space  $\mathcal{L}_p(J, R^n)$ , which contains both  $\mathcal{L}_{p_j}(J, R^n)$  and  $\mathcal{L}_2(J, R^n)$ , because  $p \geq \max\{2, p_j : j = 1, \dots, m\}$ . These operators satisfy the Lipschitz condition (4.6) and the sub-linear growth condition (4.7) as well. Therefore, Eq. (4.1) with  $F_j, G_j$  so constructed has a unique solution  $y \in \mathcal{M}_p(J)$  satisfying the initial condition  $y(0) = \varphi(0)$ . Applying Proposition 4.1 completes the proof.  $\square$

**R e m a r k 4.1.** As  $\mathcal{M}_p([0, T], R^n) \subset \mathcal{L}_p([0, T], R^n)$ , the delay operators (3.1) and (3.2) satisfy condition (3) of Theorem 4.2.

## References

- [1] J.-C. Pedjeu, G. S. Ladde, “Stochastic fractional differential equations: Modeling, method and analysis”, *Chaos, Solitons & Fractals*, **45** (2012), 279–293.
- [2] G. Jumarie, “Modified Riemann-Liouville derivative and fractional Taylor series of nondifferentiable functions further results”, *Computational Mathematics and Applications*, **51**:9-10 (2006), 1367–1376.
- [3] B. Øksendal, *Stochastic Differential Equations. An Introduction with Applications*, Springer, 2014.
- [4] I. Neveu, *Discrete Parameter Martingales*, North-Holland, Amsterdam, 1975.
- [5] N. V. Azbelev, V. P. Maksimov, L. F. Rakhmatulina, *Introduction to the Theory of Functional Differential Equations. Methods and Applications*, Hindawi, New York, 2007.

## Information about the author

**Arcady Ponosov**, Doctor of Natural Sciences, Professor of the Institute of Mathematics. Norwegian University of Life Sciences, Ås, Norway.  
E-mail: arkadi@nmbu.no  
**ORCID:** <https://orcid.org/0000-0001-5018-6577>

Received 25.01.2023  
Reviewed 28.02.2023  
Accepted for press 10.03.2023

## Информация об авторе

**Поносов Аркадий Владимирович**, доктор естественных наук, профессор Института Математики. Норвежский университет естественных наук, г. Ос, Норвегия.  
E-mail: arkadi@nmbu.no  
**ORCID:** <https://orcid.org/0000-0001-5018-6577>

Поступила в редакцию 25.01.2023 г.  
Поступила после рецензирования 28.02.2023 г.  
Принята к публикации 10.03.2023 г.